



Specimen Label

ACTIVE INGREDIENTS:

Clothianidin 3.20%

Pyriproxyfen 3.90%

OTHER INGREDIENTS 92.90%

100.00%

This product contains 0.269 lbs. Clothianidin per gallon and 0.328 lbs. Pyriproxyfen per gallon

KEEP OUT OF REACH OF CHILDREN

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information regarding medical emergencies or pesticide incidents, call 1-888-740-8712.

See inside for Precautionary statements.

KILLS:

Ants (including Red Imported Fire Ants, Harvester, Pharaoh and Carpenter Ants)	Asian Lady Beetles	Centipedes	German Cockroaches	Silverfish Sowbugs	Japanese Beetles
	Kudzu Bugs	Springtails	House Flies	Chinch Bugs	Oriental Beetles
	Brown Marmorated Stink Bugs	Crickets	Millipedes	Aphids	Boxelder Bugs
		Mole Crickets	Mosquitoes	Crane Flies	Scales
		Earwigs	Pillbugs	June Beetles	
		Fleas			

CONTROLS:

Ants (including Red Imported Fire Ants, Harvester, Pharaoh and Carpenter	Asian Lady Beetles	Stink Bugs	Pillbugs	June Beetles
	Kudzu Bugs	Springtails	Silverfish	Japanese Beetles
	Brown Marmorated	Crickets	Sowbugs	Oriental Beetles
		Mole Crickets	Chinch Bugs	Boxelder Bugs
		Earwigs	Aphids	Scales
		Millipedes	Crane Flies	

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read and follow ALL label directions when using this product.

USE RESTRICTIONS:

- NOT REGISTERED FOR SALE, DISTRIBUTION OR USE IN THE STATE OF NEW YORK.
- Do not apply to humans, animals, clothing or human bedding.
- Do not apply within aircraft cabins.
- Do not use in outdoor residential misting systems.
- Do not apply to household pets. Cover fish tanks prior to application near tanks.
- Do not apply in food/feed areas of food handling establishments.
- When applying in residential or commercial structures, do not allow spray to contact food or food-containing surfaces, feed or water supplies. Thoroughly wash dishes and food handling utensils with soap and water if they become contaminated with this product.
- Do not allow application to drip or runoff from surfaces. Do not water the treated area to the point of runoff.
- Do not apply this product in poultry houses with a mechanically-pressurized handgun.
- Not for use on plants being grown for sale or other commercial use. Not for use in nurseries, sod farms and golf courses.
- Keep all people and pets out of areas (indoors and outdoors) being treated and restrict access to these areas until all surfaces are dry.
- For outdoor applications: Do not apply more than 0.4 lbs. Clothianidin per acre per year or 1.49 gallons of Sumari™ Insecticide per acre per year or more than 4.35 fl. oz. of Sumari Insecticide per 1,000 sq. ft. per year.

Use only in areas described. For best results, follow directions for specific use areas.

USE AREAS:

For use, indoors and outdoors, in non-food areas: backyards, lawns, trees, ornamental landscaping, recreational areas, parks and athletic fields. Also for use in and around homes, schools, warehouses, office buildings, apartment buildings, theaters, hotels, resorts, industrial buildings, motels, camp sites/campgrounds, aircraft cargo areas, kennels, livestock and poultry housing, horse stables, feed lots, zoos, food processing plants, food service establishments, restaurants, outdoor dining areas, supermarkets and grocery stores, transportation equipment, truck trailers, railroad cars and food manufacturing and warehousing establishments.

This product should not stain or cause damage to painted or varnished surfaces, plastics, fabrics, or other surfaces where water applied alone causes no damage. Treat a small area and allow it to dry to determine whether staining will occur.

APPLICATION METHODS:

Outdoor broadcast spray treatment	Structural perimeter treatment
Foliar spray treatment	Surface spray treatment
Spot, crack and crevice treatment	

APPLICATION EQUIPMENT:

Including:	Backpack sprayers	Mist sprayers	Power sprayers
Backpack mist blowers	Handheld sprayers	Mechanical sprayers	

MIXING INSTRUCTIONS:

Sumari Insecticide should be diluted with water only and applied with hand-pressurized or power-operated equipment. Do not mix this product with oil. Clean equipment before and after use. Agitate the container of Sumari Insecticide before diluting by inverting it several times. For dilution, add approximately half the required water to the tank and then add the appropriate amount of Sumari Insecticide. Agitate and slowly add the remaining water. Agitate the mixture thoroughly before using and occasionally during use to ensure dispersion. Avoid excessive application.

Active Ingredient	Final % Concentration of Active Ingredient (1 fl. oz. per gallon)	Final % Concentration of Active Ingredient (2 fl. oz. per gallon)
Clothianidin	0.025%	0.05%
Pyriproxyfen	0.030%	0.06%
Total % Concentration of all Active Ingredients	0.055%	0.11%

INDOOR USE TO KILL LISTED PESTS:

Application Site	Pest	Dilution Rate In Water	Application Method(s)
Residential homes, business, office complexes, commercial, institutional and industrial buildings, multi-family residential complexes, Hospital, nursing homes, airports, and schools.	Ants (including Red Imported Fire Ants, Harvester, Pharaoh and Carpenter Ants), Asian Lady Beetles, Kudzu Bugs, Brown Marmorated Stink Bugs, Centipedes, Springtails, Crickets, Mole Crickets, Earwigs, Fleas, House Flies, Millipedes, Pillbugs, Silverfish, Sowbugs, Chinch Bugs, Aphids, Crane Flies, June Beetles, Japanese Beetles, Oriental Beetles, Boxelder Bugs, and Scales.	1 fl. oz. per gallon or 1 fl. oz. in sufficient water to cover 1,000 sq. ft.	Spot treatment Crack and crevice
	German Cockroaches, Mosquitoes	2 fl. oz. per gallon or 2 fl. oz. in sufficient water to cover 1,000 sq. ft.	Spot treatment Crack and crevice

For small volume indoor applications, see mixing chart below.

Apply as spot or crack and crevice. Low rate is equivalent to 1 fl. oz. per gallon. High rate is equivalent to 2 fl. oz. per gallon.

Volume	Low rate	High rate
Pint (16 fl. oz.)	1/8 fl. oz.	1/4 fl. oz.
Quart (32 fl. oz.)	1/4 fl. oz.	1/2 fl. oz.
Half Gallon (64 fl. oz.)	1/2 fl. oz.	1 fl. oz.
¾ Gallon (96 fl. oz.)	3/4 fl. oz.	1.5 fl. oz.

SPECIFIC USE DIRECTIONS:

Listed Crawling Insect/Pest:

Including: German Cockroaches, Centipedes, Crickets, Millipedes, Earwigs, Silverfish. Make spot and/or crack and crevice applications into harborage areas, cracks and crevices, entry points, under refrigerators, vending machines, around pipe and along baseboards. When applying to German roaches and centipedes, directly spray the pests.

Ants:

Make spot, crack and crevice and/or voids applications where ants may be foraging and/or nesting, including, around doors window frames, under sinks, around pipes, attic venting and other areas where ants may find access or nest.

Fleas:

Apply to infested areas, including rugs, floors, carpets, upholstered furniture, pet beds and resting areas, directly contacting fleas while spraying.

House Flies:

Directly spray house flies.

OUTDOOR USE:

Application Site	Pest	Dilution Rate In Water	Application Method(s)
Turfgrasses (Residential home lawns, business, office complexes, commercial, institutional, industrial, multi-family residential complexes, airports, cemeteries, parks, playgrounds, athletic fields.)	Ants (including Red Imported Fire Ants, Harvester, Pharaoh and Carpenter Ants), Asian Lady Beetles, Kudzu Bugs, Brown Marmorated Stink Bugs, Springtails, Crickets, Mole Crickets, Earwigs, Millipedes, Pillbugs, Silverfish, Sowbugs, Chinch Bugs, Aphids, Cutworms, Crane Flies, June Beetles, Japanese Beetles, Oriental Beetle, Boxelder Bugs, and Scales.	Light infestations: 1 fl. oz. per gallon or 1 fl. oz. in sufficient water to cover 1,000 sq. ft. Heavy infestations: 2 fl. oz. per gallon or 2 fl. oz. in sufficient water to cover 1,000 sq. ft.	Broadcast spray treatment, Foliar spray treatment, Spot treatment

SPECIFIC USE DIRECTIONS:

Exterior Structural and Surrounding Area for listed insects:

Treat pests directly where they harbor, rest, trail or hide. Treat structural voids around entryways including doors and windows, utility entry points, behind siding, weep holes, eaves, around lights and garbage cans.

Ants:

Make broadcast, spot and/or crack and crevice applications to any foraging trails, where ants are present. Make spot and/or crack and crevice applications to all exterior entry points around the structure. Curative and preventative broadcast applications can be made to yards, lawns, fields, parks, cemeteries, and landscaping.

For up to 90 day residual control, apply at these rates per gallon or in sufficient water to cover 1,000 sq. ft.:

Surface/Location	Fire Ants	Pharaoh Ants	Carpenter Ants
Porous surface: outdoors	2 fl. oz.	2 fl. oz.	2 fl. oz.
Porous surface: indoors	1 fl. oz.	1 fl. oz.	2 fl. oz.
Non porous surface: outdoors	2 fl. oz.	1 fl. oz.	2 fl. oz.
Non porous: indoors	1 fl. oz.	1 fl. oz.	2 fl. oz.

Fleas:

Directly spray fleas in lawns, yards, porches, patios or other areas fleas are present. Directly spray fleas in shaded areas where pets or feral animals frequent. Apply directly to fleas located in pet entry areas including steps and stoops and other areas where pets rest or sit for extended periods of time.

House Flies:

Directly spray house flies where observed on surfaces of screens, doors, window frames, and porches, in garages, on foliage or wherever flies enter or congregate around or on the building. Directly spray house flies located inside and outside of trash cans, dumpsters and other areas where garbage is kept.

PRECAUTIONARY STATEMENTS

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Mixers, loaders, applicators, and other handlers must wear long-sleeved shirt, long pants, socks and shoes.

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic organisms, including fish and invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutter or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid runoff to water bodies or drainage systems. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not apply where runoff is likely to occur.

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively foraging in the treatment area.

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

PROTECTION OF POLLINATORS



APPLICATION RESTRICTIONS EXIST FOR THIS PRODUCT BECAUSE OF RISK TO BEES AND OTHER INSECT POLLINATORS. FOLLOW APPLICATION RESTRICTIONS FOUND IN THE DIRECTIONS FOR USE TO PROTECT POLLINATORS.

Look for the bee hazard icon in the Directions for Use for each application site for specific use restrictions and instructions to protect bees and other insect pollinators.

This product can kill bees and other insect pollinators

Bees and other insect pollinators will forage on plants when they flower, shed pollen, or produce nectar.

Bees and other insect pollinators can be exposed to this pesticide from:

- o Direct contact during foliar applications, or contact with residues on plant surfaces after foliar applications
- o Ingestion of residues in nectar and pollen when the pesticide is applied as a seed treatment, soil, tree injection, as well as foliar applications.

When using this product, take steps to:

- o Minimize exposure of this product to bees and other insect pollinators when they are foraging on pollinator attractive plants around the application site.
- o Minimize drift of this product on to beehives or to off-site pollinator attractive habitat. Drift of this product onto beehives or off-site to pollinator attractive habitat can result in bee kills.

Information on protecting bees and other insect pollinators may be found at the Pesticide Environmental Stewardship website at:

<http://pesticidestewardship.org/PollinatorProtection/Pages/default.aspx>.

Pesticide incidents (for example, bee kills) should immediately be reported to the state/tribal lead agency. For contact information for your state, go to: www.aapco.org. Pesticide incidents should also be reported to the National Pesticide Information Center at: www.npic.orst.edu or directly to EPA at: beekill@epa.gov

PHYSICAL OR CHEMICAL HAZARDS

Do not apply this product around electrical equipment due to the possibility of shock hazard.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE:

Store in a cool, dry place. Always store pesticides in the original container. Store away from food and pet food.

PESTICIDE DISPOSAL:

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING:

Non-refillable container. Do not reuse or refill this container.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling if available for reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.



EPA Reg. No. 1021-2827
EPA Est. No. 1021-MN-2
Rev. 0919-0220



Manufactured for MGK®.
©2020 MGK. All rights reserved.
MGK® is a registered trademark of McLaughlin Gormley King Company
7325 Aspen Lane North, Minneapolis, MN 55428