

GROUP

4A

INSECTICIDE



Safari 2 G INSECTICIDE



For systemic insect control
in ornamental plants

For Greenhouse, Interior
Plantscape, Lath and

Shadehouses, Nursery and Outdoor
Landscape Use

Active Ingredient:	By Wt.
*Dinotefuran	2%
Other Ingredients	98%
Total	100%

*N-methyl-N'-nitro-N'-[(tetrahydro-3-furanyl)methyl]
guanidine

EPA Reg. No. 59639-149

EPA Est. No. 67545-AZ-01

KEEP OUT OF REACH OF CHILDREN

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact **800-892-0099** for emergency medical treatment information.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear: Long-sleeved shirt and long pants, chemical-resistant gloves made of any waterproof material and shoes plus socks.

USER SAFETY REQUIREMENTS

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions exist for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to shrimp. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in water adjacent to treated areas. Do not dispose of equipment washwaters or rinsate into a natural drain or water body. Do not contaminate water when disposing of equipment washwaters or rinsate.

This compound is toxic to honey bees. The persistence of residues and potential residual toxicity of dinotefuran in nectar and pollen suggest the possibility of chronic risk to honey bee larvae and the eventual instability of the hive.

This product is toxic to bees exposed to treatment for more than 38 hours following treatment. Do not apply this product to blooming, pollen-shedding or nectar-producing parts of plants during this time period, unless the application is made in response to a public health emergency declared by appropriate state and federal authorities.

Dinotefuran and its degradate, MNG, have the properties and characteristics associated with chemicals detected in ground water. The high water solubility of dinotefuran, and its degradate, MNG, coupled with its very high mobility, and resistance to biodegradation indicates that this compound has a strong potential to leach to the subsurface under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination. Periodic monitoring of shallow groundwater in the use area is recommended.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, greenhouses and handlers of agricultural insecticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is: long-sleeved shirt and long pants, chemical-resistant gloves made of any waterproof material and shoes plus socks.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

RESISTANCE MANAGEMENT RECOMMENDATIONS

Safari® 2 G Insecticide contains a Group 4A insecticide. Insect biotypes with acquired resistance to Group 4A insecticides may eventually dominate the insect population if Group 4A insecticides are used repeatedly in the same field or in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by Safari 2 G or other Group 4A insecticides.

To delay the development of insecticide resistance in greenhouse, nursery and interiorscape use sites, strongly consider the following recommendations:

- Do not apply Safari 2 G or other Group 4A insecticides to consecutive generations of the same insect pest species.
- Do not apply Safari 2 G or other Group 4A insecticides more than one time per crop cycle or three months, whichever is shorter.
- Base insecticide use on a comprehensive Integrated Pest Management (IPM) program.
- Monitor treated insect populations for loss of field efficacy.

(Continued)

RESISTANCE MANAGEMENT RECOMMENDATIONS, Continued

- Contact your local extension specialist, certified crop advisors, and/or manufacturers for insecticide resistance management (RM) and/or IPM recommendations for the specific site and resistant pest problems.
- For further information contact Valent U.S.A. Corporation at the following toll free number: 800-682-5368.

DISCLAIMER, RISKS OF USING THIS PRODUCT, LIMITED WARRANTY AND LIMITATION OF LIABILITY

IMPORTANT: Read the entire Label including this Disclaimer, Risks of Using this Product, Limited Warranty and Limitation of Liability before using this product. If the terms are not acceptable THEN DO NOT USE THE PRODUCT; rather, return the unopened product within 15 days of purchase for a refund of the purchase price.

RISKS OF USING THIS PRODUCT

The Buyer and User (referred to collectively herein as "Buyer") of this product should be aware that there are inherent unintended risks associated with the use of this product which are impossible to eliminate. These risks include, but are not limited to, injury to plants and crops to which this product is applied, lack of control of the target pests or weeds, resistance of the target pest or weeds to this product, injury caused by drift, and injury to rotational crops caused by carryover in the soil. Such risks of crop injury, non-performance, resistance or other unintended consequences are unavoidable and may result because of such factors as weather, soil conditions, disease, moisture conditions, irrigation practices, condition of the crop at the time of application, presence of other materials either applied in the tank mix with this product or prior to application of this product, cultural practices or the manner of use or application (or a combination of such factors) all of which are factors beyond the control of Valent. The Buyer should be aware that these inherent unintended risks may reduce the harvested yield of the crop in all or a portion of the treated acreage, or otherwise affect the crop such that additional care, treatment and expense are required to take the crop to harvest. If the Buyer chooses not to accept these risks, THEN THIS PRODUCT SHOULD NOT BE APPLIED. By applying this product Buyer acknowledges and accepts these inherent unintended risks AND AGREES THAT ALL SUCH RISKS ASSOCIATED WITH THE APPLICATION AND USE ARE ASSUMED BY THE BUYER.

(Continued)

RISKS OF USING THIS PRODUCT, Continued

Valent shall not be responsible for losses or damages (including, but not limited to, loss of yield, increased expenses of farming the crop or such incidental, consequential or special damages that may be claimed) resulting from use of this product in any manner not set forth on the label. Buyer assumes all risks associated with the use of this product in any manner or under conditions not specifically directed or approved on the label.

LIMITED WARRANTY

Valent warrants only that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the label, under average use conditions, when used strictly in accordance with the label and subject to the **Risks of Using This Product as described above. EXCEPT AS SET FORTH ABOVE, VALENT MAKES NO OTHER WARRANTIES, EITHER EXPRESSED OR IMPLIED.** No agent or representative of Valent or Seller is authorized to make or create any other express or implied warranty.

LIMITATION OF LIABILITY

In no event shall Valent or Seller be liable for any incidental, consequential, indirect or special damages resulting from the use or handling of this product. The limitation includes, but is not limited to, loss of yield on all or any portion of the treated acreage, increased care, treatment or other expenses required to take the crop to harvest, increased finance charges or altered finance ratings, emotional or mental distress and/or exemplary damages. **THE EXCLUSIVE REMEDY OF THE BUYER, AND THE EXCLUSIVE MAXIMUM LIABILITY OF VALENT OR SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT SHALL BE THE RETURN OF THE PURCHASE PRICE OF THIS PRODUCT OR, AT THE ELECTION OF VALENT OR SELLER, THE REPLACEMENT OF THIS PRODUCT.**

PROMPT NOTICE OF CLAIM

Valent must be provided notice as soon as Buyer has reason to believe it may have a claim, but in no event later than twenty-one days from date of planting, or twenty-one days from the date of application, whichever is latter, so that an immediate inspection of the affected property and growing crops can be made.

If Buyer does not notify Valent of any claims, in such period, it shall be barred from obtaining any remedy.

(Continued)

NO AMENDMENTS

Valent and Seller offer this product, and Buyer accepts it, subject to the foregoing **Disclaimer, Risks of Using This Product, Limited Warranty and Limitation of Liability**, which may not be modified by any oral or written agreement.

TANK MIXES

NOTICE: Tank mixing of this product with any other product which is not specifically and expressly authorized by the label shall be the exclusive risk of user, applicator and/or application advisor.

Read and follow the entire label of each product to be used in the tank mix with this product.

GENERAL INFORMATION

Applications of Safari 2 G in outdoor residential areas may be made by commercial licensed applicators only.

ORNAMENTAL PLANTS

- *Safari 2 G* can be applied as a broadcast or single plant treatment for insect control in ornamental plants in greenhouses, nurseries, outdoor landscapes and interior plantscapes.
- *Safari 2 G* is a systemic product and will be taken up by the root system and transported throughout the plant.
- *Safari 2 G* will be transported more quickly in herbaceous plants than in woody shrubs and trees. Speed of insect control will range from as little as one day for small herbaceous plants in containers, to several weeks in large trees growing in the landscape. Plant growth rate, rainfall and irrigation following application will affect speed of control.
- For nursery and landscape ornamentals, broadcast applications cannot exceed a total of 27 lbs of product (0.54 lbs active ingredient) per acre per year.

RESTRICTIONS AND LIMITATIONS

- Do not graze treated areas or use clippings from treated areas for feed or forage.
- Avoid runoff or puddling of irrigation water following application.
- Keep children and pets off treated areas until dusts have settled.
- Avoid application to areas that are water logged or saturated, or frozen, which will not allow penetration into the root zone of the plant.

APPLICATION PROCEDURES

For flower beds, broadcast *Safari 2 G* evenly over the top of plants. For treatment of individual shrubs or trees, broadcast *Safari 2 G* evenly on soil around base of plant within 18" of the trunk. For best control, pull back mulch prior to application if mulch is more than 1" thick. Irrigate after application with enough water to dissolve granules and move product into root zone.

PESTS CONTROLLED BY SAFARI 2 G	
Adelgids including: Hemlock Woolly Flatheaded Borers including: Flatheaded Appletree Fungus Gnats (larvae) Glassy-Winged Sharpshooter Lacebugs including: Azalea Leaf Beetles Leafhoppers Leafminers including: Boxwood Serpentine	Scales (Armored and Soft) including: Cottony Cushion Cryptomeria Cycad Aulacaspis Duplacionaspis Elongate Hemlock Euonymus False Oleander Fletcher Florida Red Florida Wax Oystershell Pine needle Tea Tulip Tree
(Continued)	

PESTS CONTROLLED BY SAFARI 2 G, Continued	
Mealybugs including: Citrus Long-tailed Madeira Obscure Pink Hibiscus Root Psyllids including: Asian Citrus Root Weevils (larvae and adults) including: Black Vine Roundheaded Borers including: Asian Longhorned	Thrips including: Chilli Western Flower (suppression) Whiteflies including: Giant Greenhouse Silverleaf/Sweetpotato (B and Q biotypes) White Grubs including: Oriental Beetle

RECOMMENDATIONS FOR APPLICATION TO SOIL MEDIA OF CONTAINERIZED PLANTS

For systemic insect control of containerized ornamental plants in nurseries, greenhouses, interior plantscapes, lath and shadehouses and outdoor landscapes (commercial, industrial, recreational and outdoor residential) by broadcast or individual pot application.		
CROPS	PRODUCT RATES	REMARKS
Ornamental Plants including: Bedding Plants Evergreens Flowering Plants Foliage Plants Ground Covers Non-Bearing Fruit Trees Non-Bearing Nut Trees Non-Bearing Vines Ornamental Trees Shrubs	Treatment of Individual Pots	Evenly apply granules to soil surface.
	Pot Diameter (inches)	Only apply to moist soil media. Do not apply to dry or saturated media.
	Teaspoons (tsp) per Pot	For optimal performance, do not apply until roots from transplanted plugs or liners have grown at least 1/2 inch.
	4	1/8 to 1/4 tsp
	5	1/4 to 1/3 tsp
	6	1/3 to 2/3 tsp
	7	2/3 to 3/4 tsp
	8	3/4 to 1 tsp
	For larger pot volumes, topically apply 1/3 to 2/3 tsp (1.3 to 2.6 grams) to soil surface for each gallon of potting media in container.	Irrigate pots after application with just enough water to dissolve granules.
	Broadcast Application (Pot Diameter less than 4") Apply 0.62 lbs per 1,000 sq ft of bench space or plant bed. 27 lbs/A (0.54 lb ai/A)	Do not leach treated soil media for at least 7 days after application or performance may be reduced. Heavy rainfall or excessive irrigation following application may decrease performance. In general, higher rates will be needed to control insects on woody plants than on herbaceous plants. Poinsettia: For optimal control of whiteflies, treat plants 1 to 3 weeks after pinch. Later season drenches will take longer to give control. Broadcast application may be less residual than single pot rates, especially on woody plants.
<ul style="list-style-type: none"> One level teaspoon contains 4.0 grams and 1 cup (8 fl oz) contains 6.6 oz of Safari 2 G. Refer to "Resistance Management Recommendations" section of the label for additional guidelines. Do not apply more than 27 lbs (0.54 lbs ai) per acre of nursery or landscape per year. 		

RECOMMENDATIONS FOR SOIL APPLICATION TO IN-GROUND PLANTS

For systemic insect control on ornamental plants in field nurseries (in-ground), interior plantscapes and outdoor landscapes (commercial, industrial, recreational and outdoor residential) by broadcast application, or as an individual plant treatment.

CROPS	PRODUCT RATES	REMARKS
Ornamental Plants including: Bedding Plants Evergreens Flowering Plants Foliage Plants Ground Covers Non-Bearing Fruit Trees Non-Bearing Nut Trees Non-Bearing Vines Ornamental Trees Shrubs	Treatment of Individual plants Trees 2.0 to 4.0 oz per inch of trunk diameter at breast height (DBH) For multi-stem trees, base rate on cumulative inches of diameter of all stems at breast height.	For optimal control, apply early in plant's annual growing cycle and keep soil moist for at least 7 days after application. Only apply to moist soil. Do not apply to dry, saturated or frozen soil or when plants are not actively taking up water from soil. Heavy rainfall or inadequate irrigation immediately following application may decrease performance.
	Shrubs 1.0 to 2.0 oz per foot of shrub height	Use higher labeled rates for very diameter trees and for broadleaf evergreens with dense foliage (example hollies).
	Broadcast application in herbaceous flower beds and ground covers Apply 0.62 lb per 1,000 sq ft of bed 27 lbs/A (0.54 lb ai/A)	Broadcast application may not be effective in shrubs and trees.
<ul style="list-style-type: none"> One level teaspoon contains 4.0 grams and 1 cup (8 fl oz) contains 6.6 oz of <i>Safari 2 G</i>. Do not apply more than 27 lbs (0.54 lbs ai) per acre of nursery or landscape per year. To delay the development of resistance, do not make more than one soil application per crop cycle or three months, whichever is shorter. Refer to "Resistance Management Recommendations" section of the label for additional guidelines. 		

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.

PESTICIDE STORAGE

Keep pesticide in original container.

Do not put concentrate or dilute into food or drink containers.

Store in a cool, dry place.

Do not store diluted spray.

For help with any spill, leak, fire or exposure involving this material, call day or night 800-892-0099.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Triple rinse (or equivalent). Do not reuse container.

Offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Manufactured for:

Valent U.S.A. Corporation

P. O. Box 8025

Walnut Creek, CA 94596-8025

Made in U.S.A.

Form 1671-A

EPA Reg. No. 59639-149

EPA Est. No. 67545-AZ-01

Information contained in this booklet is accurate at the time of printing. Since product testing is a continuous process, please read and follow the directions on the product label for the most current directions and precautionary statements.

Always check with your state to verify state registration status.



For state registration and/or supplemental labels, please call or visit us online.

Products That Work, From People Who Care® | www.valent.com | 800-89-VALENT (898-2536)

Read and follow the label instructions before using.

Safari and *Products That Work, From People Who Care* are trademarks of Valent U.S.A. Corporation. ©2008 Valent U.S.A. Corporation. All rights reserved. Printed in the USA. 2008-SAF2G-0001 2/08 mfl/AV